

## AMENDMENTS TO THE SPECIFICATION

Please amend page 12, line 26 as follows:

Next, ~~12~~ 1.1 g of Coenzyme Q10 are dissolved into the mixture.

Please amend page 48, lines 23-25 as follows:

A topical formulation of Coenzyme Q10 (~~10%~~ A 1.0% and 1.5% formulation was tested) was applied to the tumors in the treatment group daily for a period of 30 days.

Please amend page 49, lines 5-10, as follows:

~~6g~~ 10g of Phospholipon 90G and 5g of Phospholipon 90H (American Lechitin, Stanford, CT) was dissolved in a mixture of ~~5.8g~~ 13.3g of Glycerol (Sigma-Aldrich, St. Louis, MO), ~~0.2g~~ 0.3g BHT (Sigma-Aldrich), ~~4ml~~ 9ml ethanol (Sigma-Aldrich), and ~~18g~~ 1.5g MCT (Sigma-Aldrich) at 60°C. ~~20g~~ 1.1g of CoQ10 (Pure Prescriptions) were dissolved into the resulting mixture. ~~90ml~~ 65ml of 1mM phosphate buffer (pH 8.2) prepared with nitrogen saturated water and 0.2ml of lavender (Sigma-Aldrich) were added and the mixture was blended in a high speed blender at 12,000 RPM to form a cream. The cream was stored at 4°C until used.